Onsite Dust Monitoring

4/1/2020 – 4/30/2020

Net Dust (All Downwind Stations)



Net dust represents the dust that may be leaving the Site. This is determined by subtracting upwind data (dust blowing onto the Site from other sources) from downwind data. This helps us monitor that dust control actions are effective. No data was recorded over the Easter Holiday (April 9-12).

Offsite Dust Monitoring

Total dust readings including upwind dust contribution

Weekly - 4/17/2020 – 4/23/2020

Individual Offsite Stations:
24-Hr Average Dust Readings (µg/m³)











Notes: California Ambient Air Quality Standard for PM10 averaged over 24 hours is 50 µg/m³. National Ambient Air Quality Standard for PM10 averaged over 24 hours is 150 µg/m³.



South Coast Basin Regional PM10:
24-Hr Average Readings (µg/m³)



Closest regional station provided for comparison to regional trends

24-average concentrations were below air quality standards. Winds were blowing primarily from the west, and to a lesser extent from the west/southwest.

Onsite Dust Monitoring

Total dust readings including upwind dust contribution

Weekly - 4/17/2020 – 4/23/2020

Individual Onsite Stations:
24-Hr* Average Dust Readings (µg/m³)



* Note: DT-1, DT-3 and DT-5 represent the average dust readings for working hours only.



24-average concentrations were below air quality standards. Winds were blowing primarily from the west, and to a lesser extent from the west/southwest.

Notes: California Ambient Air Quality Standard for PM10 averaged over 24 hours is 50 µg/m³. National Ambient Air Quality Standard for PM10 averaged over 24 hours is 150 µg/m³.

Closest regional station provided for comparison to regional trends.

South Coast Basin Regional PM10:
24-Hr Average Readings (µg/m³)